

ES 10 March 2, 2001
Bartolome

Soil development

1. Origins of soil science
2. General characteristics of soils
3. Jenny's state factor equation $s=f(Cl,o,r,p,t)$
4. Soil description and classification

Figures: 3.10 3.13 Allaby; 3.14 Allaby; 4.14 Schlesinger




Figure 3.13 Profiles of four soils, with the vegetation associated with them

The 11 soil orders of the US soil taxonomy

Alfisols	Soils of climates with 510–1270 mm annual rainfall; most develop under forests; clay accumulates in the B horizon.
Andisols	Volcanic soils, deep and light in texture; contain iron and aluminium compounds.
Aridisols	Desert soils with accumulations of lime or gypsum; often with salt layers; little organic matter.
Entisols	Little or no horizon development; often found in recent flood plains, under recent volcanic ash, as wind-blown sand.
Histosols	Organic soils; found in bogs and swamps.
Inceptisols	Young soils; horizons starting to develop; often wet conditions.
Mollisols	Very dark soils; upper layers rich in organic matter; form mainly under grassland.
Oxisols	Deeply weathered soils; acid; low fertility; contain clays of iron and aluminium oxides.
Spodosols	Sandy soils found in forests, mainly coniferous; organic matter, iron and aluminium oxides accumulated in B horizon; strongly acid.
Ultisols	Deeply weathered tropical and subtropical soils; strongly acid; clay accumulated in B horizon.
Vertisols	Clay soils that swell when wet; develop in climates with pronounced wet and dry seasons; deep cracks appear when dry.




Figure 4.14 Annual sediment flux from major drainage basins to the world's oceans. Data are millions of tons (10^{12} g) per year, and arrows are drawn proportional to the flux. From Milliman and Meade (1983).




Figure 3.14 World distribution of soil orders